



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

566
.Y6
J66

B 909,978

HANGING VALLEYS OF THE YOSEMITE

revised by
D. W. JOHNSON
Harvard University

REPRINTED FROM THE
BULLETIN OF THE AMERICAN GEOGRAPHICAL SOCIETY
VOL. XLIII., NOVEMBER-DECEMBER, 1911

1

1

the Hanging Valley peculiar notches w valleys and has four and not ice was ma short criticism of th Yosemite Valley, is quite an ordinary an other papers publish of "Glacial Erosion W. P. Blake (1900) the author maintains the "Profile of Matun (1904) regards abno result of glacial deep by Gravity Streams w (1909) treats the Yos

The following page River in the Yosemite tributary streams; the Merced, by which the cess of valley widening present width of the of the hanging valley and their relation to

THE FORMER

In an earlier number set forth in some detail tributary hanging valleys with reasonable accuracy main valley has suffered discussion of the limitations of measurements of gradient courses of tributaries, the streams having highly irregular measurement streams. If several miles in tributaries to their profiles o main stream may be asce danger of error being large

the amount of over-
is, according to the
2,575 feet, — the deepening of the
moved through it (I believe that the valley
glacial erosion; but may be such as material
deepening.

The Valley Map of
Creek and of the Eastern
short streams, but Canyon both appear
level of the Merced
than a mile above the elevation of the Merced
similar portion of Eastern
level of the Merced.

Snow Creek is a branch of
Valley Map. The grade is
over-steepened lower
feet as the former elevation
it emptied. The average
Snow Creek, as shown, is
somewhat higher elevation
branch.

The Valley Map represents
Creek, and shows it to be in
relations as portrayed on
that the lower course of
and that the Bridal Veil
discordance in the junction
measurements on five or six
ungraded lower portion,
mate former elevation of
Illionette Creek presented
mentioned above. Only
are the Valley Map and for the
four in the over-steepened
the the quadrangle, the
the position of the

in the evidence
elevation of the
graded. In plot
I have tried to es-
mit sufficient acc
narrow limits, wh
to say, however,
dence that the Yo
tent of from 2,200
of the valley, and co
of the valley floor in
glacial trough, we a
from 300 to possibly
feet or more. A mi
servative way of stat
of this profound over-

THE

In concluding his
Turner mentions four
for the formation of

1. That it was
2. That it was
3. That it was
4. That it was

due to the sapping
Conte? and Russ
jointing (Becker
Branner,

To the objections urged
the above theories (faulting,
ing valleys were evidently gra
and Tenaya Creek before t
formed; the relation of trib
changed by the formation of t
their hanging character; hence
the down-faulted block must
irregular course of the establi
sent off a branch graben at it
already established Tenaya Cree
able that we must look to norm
for an explanation of the Yosemite

merced River; and
s largely respons-

during the Cre-
ied by the Sierra
face of relatively
ivers. Many of
als by lava flows
The close of the
res of the pres-
end of the Neo-
reatly increased
id caused them
ie striking ele-
on cutting was
ich ultimately
some distance

ned the back
hich the can-
antage from
gh it would
nearly base-
of the Pied-
pland areas
one is very
erpretation
ed and dis-
ulted from
istributive
the pres-
er's argu-
streams to
as, seems
s on the
he grade
l be true
ed block
plied to
ave the
otective
deeper

canyon cutting farther down the slope seems unnece-
incompatible with the widespread evidence of gla-
stead of protection, in the glaciated areas.

Consequent upon the uplifting and tilting of the
leveled surface, the Merced and other large so-
streams must have entrenched themselves very rapidly
tributaries deepened their valleys more gradually, es-
flowed at right angles to the main stream and hence
their gradients affected by the southwest tilting. The
valleys would come to hang above the main stream.
ment of the hanging relation was presumably most pro-
the middle courses of the Sierran rivers; for near the
hills the amount of uplift is so slight that no great
junction of main and tributary streams is possible;
headwater areas the contrast in volume between the m-
a tributary decreases to such an extent that one is al-
ful as the other. Even if tilting increased the gradie-
stream alone, in the headwater areas, there could have
great a discordance in junction as would occur far
valley where the main stream not only experienced the
in gradient, but was also a much more powerful st-
tributaries.

There is a distinct limit to the time during which ha-
can remain as such. As soon as a main stream has e-
trenched itself, and begins to establish a profile of gentle
energy decreases. The tributary streams continue to
valleys until accordant junctions with the main stream
lished. It is a well-substantiated law of stream action
required for a main stream to reduce its channel to a f-
and to open out its valley floor to a width not many t-
than the width of the stream, is longer than the time
hanging tributaries to reduce their valleys to accorda-
Accordingly, while hanging valleys might exist along
and similar rivers, especially along the middle courses of
so long as the main streams occupied narrow V-shape
fairly steep gradient, it is safe to say that the more open
main valley floor became, the less would the tributary v-
and that where the main valley floor had come to be s-
wider than the stream, and of very low gradient, no han-
would exist.

If we proceed on the assumption that the glaciers wh-
the upper portions of some of the Sierra canyons we

directly opposed to
the glaciated areas,
it exist, as before.
valleys, or hang-
her down stream,
ortant element in
sive glacial over-
And since an
wider than the
ven reverse the
old expect, after
alleys associated
former glacier)
bris plain over
will.
Sierra region
ure of Sierran
tly, the uplift
icient to cause,
many of the
persists in a
ies are much
tribution and
nificant. In
ion-glaciated
t if they are
ssive in the
er east; that
s difficult to
f glaciation
have there
ith the still
the lower
ing of the
the creek
ists. The
ra Quad-
urse; the
y would,
or. Bull
along its

course which would intersect the North Fork, to w
tributary, something less than 400' above the pres
Fork. It is apparent that these hanging valleys, wh
not typical, and whose discordance in junction is us
hundred feet, are not to be compared with the hangin
Yosemite, where tributary streams cascade abruptly




FIG. 2—Yosemite Valley looking east.

valley walls, and have a quite uniform discordance in
more than two thousand feet.
. The hanging valleys west of El Portal enter a narrow
gorge of fairly pronounced gradient. The hanging va
Yosemite enter a main valley (Figure 2), whose floor is
the width of the main stream, and whose gradient is un

wide open valleys, such as the Yosemite, which appear to afford strong evidence in favor of glacial erosion.

Instead of disproving the glacial origin of the Yosemite, the comparison of valleys suggested by Turner really affords strong evidence in favor of the theory of glacial erosion. It is no mere coincidence that of the six valleys named by that writer the two most unlike the Yosemite have presumably suffered little or no glaciation, whereas the four most like the Yosemite served as the outlets of extensive glacial basins. The areas of these basins may be determined roughly by drawing boundaries which shall enclose all of the strongly glaciated region tributary to the several valleys. Although this does not take account of the passage of ice from one basin to another across the present divides, the method is sufficiently accurate, when applied to alpine regions of strong relief, to enable one to discriminate between profoundly and slightly glaciated basins, and between basins which favor extensive erosion of the master valley, and those which do not.

The valley near Sugar Loaf, on the South Fork of the American River, is so little like the Yosemite that the contour map leaves one in doubt as to why they should be compared; for the contours represent a V-shaped gorge without hanging valleys, although a few of the tributaries have their lower courses somewhat over-steepened. The basis of comparison with Sawmill Canyon is simply that the latter "under favorable circumstances would widen out into a Yosemite"; but since this questionable statement involves the very point at issue, the comparison is not permissible. On the other hand, the small valley on the Middle Fork of Stanislaus River at the mouth of Niagara Creek bears some resemblance to the Yosemite by reason of its steep walls and flat floor; but the contour map shows such marked evidences of glaciation in the headwater portions of streams draining into this valley that it seems quite probable that a glacier traversed the valley and gave to it its peculiar form. The area of the glacial basin is smaller in this case than in the others mentioned below, and this may account for the small size of the glacial trough on the Middle Stanislaus. Hetch Hetchy Valley is truly another Yosemite, larger than the valley on the Middle Stanislaus, because it is the outlet of a larger and more profoundly glaciated basin. It is smaller than the real Yosemite, because its glacial basin, while very large, is very asymmetrical (Fig. 3 a). The ice streams, instead of flowing comparatively short distances to the center of the basin and there uniting to form a large and powerful glacier, as in

The case of the Yosemite
almost clear across
which they joined
concentrating at one
of the ice of the




FIG. 3—Sketch maps
drainage basins.
U. S. Topogra
present di
Valle

Yosemite basin
the Hetch Hetchy
less profound &
Tehipite Va
which glaciatio

The Hanging Valleys of the Yosemite

trough form is marked, this valley is less striking than the Yosemite presumably because its glaciated basin is less symmetrical and permitted less concentration of the ice streams than was the case with its more fortunate neighbor. The Grand Canyon of Kings River is the last of the valleys mentioned by Turner, is the outlet of a splendid glacial basin (Fig. 3 e); consequently it is itself a splendid glacial trough. One might add to the above list Kern Canyon, which has the trough form well developed, because it is the outlet of a glacial basin (Fig. 3 d) in which glaciation was profound, and is so shallow as to admit of a fair concentration of ice streams. Yosemite Valley owes its preëminence to the favorable combination of several elements: (1) a remarkably symmetrical basin, which was (2) profoundly glaciated, and which permitted (3) intense concentration of ice streams, one of which was augmented by (4) overflow ice from a neighboring basin. Variations in the country rock doubtless play their part in giving to the different troughs different degrees of perfection; but I regard rock differences as of less importance than differences in the size, shape, and extent of glaciation, of the basins for which the troughs serve as outlets.

Turner has suggested (p. 318) that the Yosemite Valley may have acquired a depth of 1,000 feet by the beginning of the Pleistocene, and that this "lessens the difficulty of accounting for the present depth of the Yosemite and of other canyons of the southern Sierra." The elevation of the Merced River previous to glacial overdeepening was, as shown above, from 2,000 to 2,500 feet above the present level of the stream, or between 6,000 and 6,500 above present level. This may have been 1,000 feet below the higher parts of an imperfectly developed peneplain. In any case, the present hanging valleys appear to have been graded with reference to that elevation of the Merced, and the indications are that the Merced at that time occupied an open valley as did the tributaries. The difficulty which confronts the theory of stream erosion of the Yosemite is not the absolute depth of the canyon, but rather the relation of the broad-floored main valley to the hanging valleys. This difficulty is lessened by imagining an earlier or later date for the initiation of canyon cutting.

The character of Tenaya Canyon has always been a serious difficulty in interpreting the Yosemite as a stream carved valley. Tenaya Creek, with a drainage area little if any larger than that of Yosemite Creek, flows through a wide open valley to an accordant junction with the aggraded floor of the open Yosemite Valley, instead of dropping 2,000 feet or more from a hanging valley. Assuming n

aining into
mining the

3 c) in
the glacial

The Hanging Valleys of the

mal stream erosion, it is hard to see why it be able to reduce its valley to grade and to a considerable width, while a similar tributary has a small beginning on so vast a work. but a suggested that the Lyell Fork of the Tuolumne River. This would account for the former headwaters of Tenaya Creek by the former volume of Tenaya Creek, under which canyon would seem less remarkable. Tur this very possibility, but on the evidence of river ages it is formerly came through Tenaya Canyon. It ties in explaining the relation of the canyon to the other hand, if we accept the very recent glacial period much ice from the divide into the Tenaya Basin, and (300-5-306) and Gannett (87) that the Tenaya stream which entered the Yosemite Valley served. The Tenaya Canyon owes its U-shaped depth below the hanging valleys to glacial erosion, an adequate explanation. The small Tenaya Creek is even more glacial, because it was carved by the main Merced River. The hanging portions of the main Merced River valley, because the Tenaya Glacier was larger than the Little Yosemite Glacier.

THE LONGITUDINAL PROFILE OF THE

Reference has been made above to the fi
streams entering the Merced west of El Por
Creek and the North Fork of the Merced
courses. They join the Merced with acc
short distance back from the main river show
while their upper courses are again of gent
be surprised to find a comparable feature in
further up stream. For while an even tilting we
gradient of the main river uniformly thu
greater volume would allow the lower course to
rapidly. If to difference in volume we add diffe
ance, it is easy to see how one part of the strear

rock might intrench itself more rapidly than a part up stream on more resistant rock, thus developing a locally steep gradient which would gradually retreat up stream. Something of this nature is shown in the longitudinal profile above (east of) El Portal. It should be remembered that, west of El Portal, the valley of the Merced is unglaciated, while east of the western end of the Yosemite, the valley has been profoundly glaciated. Between these two points, El Portal and the western end of the Yosemite, the valley has been traversed by glaciers to some extent, and shows the effects of glacial erosion very distinctly near the Yosemite, but to a decreasing extent toward El Portal. Had the pre-glacial Merced possessed a well-graded profile in this region, we should expect to find now, as a consequence of glacial erosion, a gradual decrease in gradient above El Portal, until the gradient finally became approximately flat, or even reversed, in the Yosemite Valley, where glaciation was most profound. Instead, we find a very pronounced steepening of the gradient above El Portal, similar to the steepening observed in the lower courses of the non-glaciated tributary streams farther west. It is evident that before glaciation the main river had not completely graded its course, although it had pushed the over-steepened zone as far up stream as El Portal. From this point the profile rose rapidly to a level some 2,000 feet above the main portion of the Yosemite Valley, as shown by the position of the hanging valleys. The western part of the Merced was then entrenched, the eastern part still flowing on the uplands. Glacial erosion has greatly altered the profile of the eastern part, but some trace of the pre-glacial conditions persists in the over-steepened profile east of El Portal.

VALLEY WIDENING IN JOINTED ROCKS

One of the most evident relations in the present topography of the Yosemite Valley is the control of joint planes upon weathering and erosion. Impressed by the remarkable form of the Yosemite Valley as a whole and by the important influence of jointing upon the form of the valley walls, some observers have genetically connected the two, seeking to explain the former by the latter. Unfortunately, no one who supports this view has set forth in detail just how stream erosion on jointed rocks can develop valleys essentially different from those developed by stream erosion on unjointed rocks. A brief analysis of this phase of the Yosemite problem may not be amiss. Let us first consider the process of valley development in massive, unjointed rocks.

RIVER

Some of the
for example,
er-steepened
nctions, but
use in grad-
One should
river itself;
ly increase
its course,
itself more
ock resist-
in weaker

The Hanging Valleys of the

If a main stream incises itself with such weather back but slightly before a deep treacher come to flow in a narrow chasm with more walls. The width of such a chasm is little slowly, and the stream. Tributaries may erode weathering so give rise to lateral hanging treat as the stream cuts downward. Two the higher portions of the walls retreat farther. First, the higher portions are exposed as it is in a longer time. Second, a greater weathering must proceed very near the bottom. Just at the stream level no weathering or no chance to remove the rock from above the stream level a narrow segment of rock fall down into the stream as fast as its weathering. Here the valley may be widened so weathering alone. The corresponding segment of rock may weather to a width of a mile or more. The width of the stream itself. The result above is perhaps an over elaborate suggestion in the V-shaped cross-profile of most young valleys. It should not be inferred that the young stream cuts itself while entrenching itself, for lateral erosion weathering in producing the width of the valley of a stream, when down-cutting is active, it is fair to say that during youth, valley deepening is ordinarily obscured by the far greater cutting, valley widening largely due to it. It follows that a young stream cannot develop a flat floor. Beginning with maturity, valley deepening is of importance, and valley widening may be due more to weathering. A flat valley floor, many times the width of the stream, may be developed.

Let us now consider whether the above principles apply to the case of the jointed mountains. If the valley walls are intersected by joint planes, these planes may produce the greatest effect upon the stream. They may be neither too widely nor too closely spaced.

modified in
order that
firms, they
too widely

modified in
order that
they, they
too widely

Returning to the case of the Yosemite, the undoubted characteristic of youth in the width of the main valley cannot be lost. The valley floor is broad and flat. Jointed granite weathering, are present in the valley influencing the weathering now in process. The hanging valleys confirm the conclusion. The Yosemite is not due to weathering; for the valley wall are worn back, by erosion at a faster rate than are the interstream areas. active. If the walls of the Yosemite had

tributary streams, and especially the far back from the main valley as to be effect. Instead, we find the tributary walls, or breaking away descending in slight the steep valley walls.

One can conceive that valley lateral cutting by a shifting stream a flat-floored, steep-walled valley to a comparatively negligible valley one level will widen stream walls. Granted the favorably in comparison with the slowly stream eroding laterally with traversed by vertical planes of faces the valley walls to take part in the retreating cliffs not that the broad floor and valley have this origin is evident. The latter would be forced River could have the existence on the same of young streams, and streams, demands a special an. There is a special exception to the very weak rock, between valley floor in the valley third rock, advanced the combination of hanging from their almost erosion, even where rock exist. The assistance exist valley and the will not explain badly join the width and quickly rem

the present walls. A far more reasonable hypothesis is to regard the Yosemite Valley as one of the many open main valleys with hanging tributaries which characterize every region of Alpine glaciation, and which are reasonably interpreted as the product of glacial over-deepening of pre-existing river valleys.

It may be pointed out that the remarkable effects of jointing observed in the walls of the Yosemite are indirectly due to glacial over-deepening and over-steepening. Not until glacial erosion had made the steep-sided trough, was full opportunity given for the joint blocks to fall from place leaving the angular re-entrants and vertical faces which form such an important element in Yosemite scenery. Weathering and gravity have worked to great advantage in the recent past, and the resultant features are bold and striking. In the future, as the walls wear back to more gentle slopes and the more stable position of the joint blocks causes them to waste gradually away instead of falling in large masses from the cliffs, the valley walls will become cloaked with debris, the bold features of today will give place to more flowing, graded profiles, and the grandeur of Cathedral Rocks, Three Brothers and Half Dome will be a thing of the past. The influence of jointing on weathering is thus to be interpreted as an indirect effect of the glacial origin of the valley, rather than as an important agent in the formation of the valley.

THE NOTCHES BESIDE THE FALLS

As an observer stands in the Yosemite Valley and views the cataract of the Yosemite, Lower Yosemite, Illilouette, Vernal or Nevada Falls, he notes that beside each fall is a deep notch which the stream curiously avoids, to fall over a sheer cliff, as if to provide for man the most stupendous spectacle possible. Branner has given us a good description of this peculiar topographic feature (547-553), and has attributed the formation of the notches to the action of the several streams during a time when they were displaced from their normal positions by ice. A glacier occupied the bottom of each valley, forcing the stream to take a course some distance to one side, where it was held between the ice and the valley wall. While in this position the notch was cut. When the glacier disappeared the stream returned to its former position, leaving the notch deserted. Inasmuch as the stream cut the notch back faster than the glacier eroded its channel, it was argued that glaciers must have very little eroding power, and that the Yosemite Valley must therefore be the result of normal stream erosion in jointed rocks.

The Hanging Valleys of

The present visitor to the Yosemite must find the present topographic map, which was not available at the time of his account of the notches. An inspection of the field evidence confirms the suggestion that some modification of this Lower Yosemite Falls, Illilouette Falls, up at a higher level in the floors of the cut, as required by Branner's theory. no signs of abandoned stream channels, signs of stream erosion. On the other hand, to result from the normal weathering




Fig. 6.—Yosemite Falls. Showing abandoned stream channel to the west, through which the trail passes. Vernal Fall is a notch due to weathering at

valleys would leave oblique scar
intersected the axes of the valley
sharp meeting the valley wall on
a notch as we find at each of
notches at Vernal Falls (Fig. 6):
developed on NW-SE joints
which gives form to the sc
Liberty Cap. The notch at Ill
E-SW joint plane, similar in
Nevada Falls and on the steep f
The notch at Nevada Falls (F
notch, but it does not open on
sent in Branner's sketch map

Falls do open
 which they were
 the notches are
 Notches are no
 ches are seen
 along oblique
 planes. When
 member that
 deepening of
 in valley left
 deep valley
 and likewise
 steps in the
 valley) in an
 le condition,
 sy to realize
 the granite
 break down
 wherever
 o blocks by
 ; and how
 ve blocks
 from the
 hanging
 int planes
 An oblique
 would give
 stion. The
 ls (Fig. 4)
 stem as the
 derick and
 s due to a
 appear at

am carved
 as repre-
 rely in the

glacial displacement to get the stream in
 stream flows naturally into it even now.
 that the notch was cut in the bottom of the
 ditions, and that a slight displacement,
 turned the stream into its present course

The notch beside the Upper
 Yosemite Falls is similar in
 origin to that beside the Ne-
 vada Falls. Yosemite Creek
 formerly flowed through the
 notch, but was shifted to its
 present position, doubtless by
 a late advance of the Yosem-
 ite Creek Glacier. It would
 seem that Yosemite Creek ac-
 quired its abnormal position
 earlier than did the Merced
 River at Nevada Falls, since
 the creek has had time to
 entrench itself to a noticeable extent, al-
 though the course of the creek may still be observed
 in the notch which is followed by the trail for some
 distance.

The existence of two distinct types
 in the Yosemite region is clearly shown
 in the above figures which are reproduced.
 The dual character of their
 formation has been pointed out in
 this paper had been prepared for presenta-
 tions in Boston during the last convoca-
 tion of the Matthes recognized the dual character of
 the work.

Returning to the question as to what
 concerning the relative efficiency of stream
 erosion should be noted that the notches at Lower
 Yosemite Falls and Vernal Falls afford no evidence
 that they were formed independently of stream
 erosion. Upper Yosemite Falls and Nevada Falls
 were formed after the main valley had been ov-



FIG. 5—III
 to w

The Hanging Valleys

ice advance to which displaced the stream. According to the theory of glacial erosion, deepened by ice action. Hence the notch is younger than (part of) the glacial deepening. The glacial displacement of the Post-glacial cutting has formed on the streams now pass over the ledges.



FIG. 6—Vernal Falls and Nevada Falls. Just so to weathering along joint planes, while not carved by the stream. Trails pass through

difference in volume and load of stream would appear from the relative sizes of the notches. The interpretation has been short as compared with the efficiency of glacial erosion.

CONCLUSIONS

The Yosemite Valley is a young glacial trough whose walls of jointed granite are in a position of the trough was determined by the shape of the glacier which guided the advancing glacier. The valley which guided the advancing glacier was favored by the shape of the glacier. The valley which guided the advancing glacier was favored by the shape of the glacier.

present positions.
valley was over-
ter-glacial age—
main valley, and
to their present

notches where
all lowances for



glacial time, it
post-glacial
e. On this
or the other

great depth,
weathering,
in stream.
sting river
glacial erosion
which per-
l form the

unusually vigorous Yosemite Glacier. Glacial over-deepening exceeded 2,000 feet in the Yosemite region, and produced the remarkable hanging valleys for which the region is famous. Glacial over-steepening produced the steep valley walls, and made possible the effective weathering along joint planes, to which the details of cliff sculpture are due. Many of the notches in the lips of the hanging valleys are due to the joint-controlled weathering; but some are stream carved notches, deserted by their streams because of glacial interference. Both types of notches are expectable features in the side of a glacial trough located in jointed rocks.

REFERENCES.

- ANDREWS, E. C. "Corrasion by Gravity Streams with Applications of the Ice Flood Hypothesis." *Roy. Soc. N. S. Wales, Jour. and Proc.*, vol. XLIII, pp. 204-330, 1909.
- BECKER, G. F. "The Structure of a Portion of the Sierra Nevada of California." *Bull. G. S. A.*, vol. II, pp. 49-74, 1891.
- BLAKE, W. P. "Glacial Erosion and the Origin of the Yosemite Valley." *Am. Inst. Mg. Engrs., Trans.*, vol. XXIX, pp. 823-835, 1900.
- BRANNER, J. C. "A Topographic Feature of the Hanging Valleys of the Yosemite." *Jour. Geol.*, vol. XI, pp. 547-553, 1903.
- GANNETT, HENRY. "The Origin of the Yosemite Valley." *Nat. Geog. Mag.*, vol. XII, pp. 86-87, 1901.
- JOHNSON, D. W. "Hanging Valleys." *Bull. Am. Geog. Soc.*, vol. XLI, pp. 665-683, 1909.
- JOHNSON, W. D. "Profile of Maturity in Alpine Glacial Erosion." *Jour. Geol.*, vol. XII, pp. 569-578, 1904.
- TURNER, H. W. "The Pleistocene Geology of the South-Central Sierra Nevada, with Especial Reference to the Origin of Yosemite Valley." *Cal. Acad. Sci., Proc.*, 3rd Ser., Geol., vol. I, pp. 261-321, 1900.

THE UNIVERSITY OF MICHIGAN
GRADUATE LIBRARY

DATE DUE

~~APR 6 1974~~

~~MAR 23 1974~~

~~DEC 2 1973~~

~~NOV 12 1975~~

